

Product Datasheet

Histone H3 (mono methyl R17) Antibody



Catalog No: CY5221

Reactivity: Human, Mouse ,Rat

www.abways.com

Isotype: Rabbit IgG

Applications: WB, IHC, IF/ICC

Information

UniProt ID: P68431

All Names: H3 histone family, member A; H3/A; H31; H3FA; H3FB; H3FC; H3FD; H3FF; H3FH; H3FI; H3FJ; H3FK; H3FL; HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J; histone 1, H3a; histone cluster 1, H3a; Histone H3.1;

Form: Liquid

Storage instructions: Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Storage buffer: pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purity: Affinity-chromatography

Immunogen: A synthesized peptide

Molecular Wt.: 15 kDa

Application


WB: 1:1000~1:2000

IHC: 1:100~1:250

ICC/IF: 1:100~1:250

Background

H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers.



Western blot analysis of Histone H3 (mono methyl R17) expression in HeLa whole cell lysate.

For Research Use Only. Not For Use In Diagnostic Procedures.

www.abways.com