## **Product Datasheet**

# Acetyl-Histone H3 (K14) Antibody

Catalog No: CY5272 Reactivity: Human, Rat

Isotype: Rabbit IgG Applications: WB IHC ICC/IF IP



www.abways.com

#### Information

UniProt ID: P68431

**All Names:** H3 histone family, member A; H3/A; H31; H3FA; H3FB; H3FC; H3FD; H3FF; H3FH; H3FI; H3FJ; H3FK; H3FL; HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J; histone 1, H3a; histone cluster 1, H3a; Histone H3.1;

Form: Liquid

Storage instructions: Store at +4° C short term. Store at -20° C long term. Avoid freeze / thaw cycle.

**Storage buffer:** pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

**Purity:** Affinity-chromatography **Immunogen:** Synthesized peptide

Molecular Wt.: 15kDa


### Application

WB: 1:500~1:2000 IHC: 1:50~1:200 ICC/IF: 1:50~1:200

IP: 1:30

#### Background

H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers.



Western blot analysis of extracts from C6 cells, using Acetyl-Histone H3 (K14) antibody.

For Research Use Only. Not For Use In Diagnostic Procedures. www.abways.com